

## MEDICATION LIST

Discontinue the use of any medications that may affect the outcome of allergy testing as described on the previous page. Drugs which may affect skin testing, include ANTIHISTAMINES, (including allergy, cold & sinus medicine, both prescription and non-prescription), and certain ANTIDEPRESSANTS.

The ANTIHISTAMINES listed below should be discontinued for **4 DAYS** before skin testing. The ANTIDEPRESSANTS listed should be discontinued for **1 DAY** prior to skin testing. (If you have any questions regarding medicines you are taking that are not listed, please call the office):

Allegra(Fexofenadine)	Coricidin	Rescon
Allegra-D	DA Chewable	Rescon Jr.
Alavert	DA II	Robitussin Night Time Cold
Alavert-D	Deconamine	Ryna
Actifed	Dimetapp	Rynatan
Alka-Seltzer Plus/Cold	Doxepin	Rynatuss
Allerest	Dramamine	Semprex-D
BC Allergy Sinus	Drixoral	Sine-Off Night Time
Bonine	Dura-Vent DA	Sinequan
Bromfed	Excedrin PM	Sinutab
Children's Tylenol Allergy-D	Extendryl	Sudafed Cold & Allergy
Children's Tylenol Multisymptom Cold, Chewable & Liquid	Fedahist	Tavist(Clemastine)
Chlortrimeton	Kronofed	Tavist-D
Chlorpheniramine Maleate	Librium	Thera-Flu
Clarinex	Nolahist	Triaminic
Clarinex D	Nyquil	Tussi-12
Claritin(Loratadine)	PBZ-SR	Tussionex
Claritin-D	PediaCare	Tylenol Allergy Sinus
Comtrex	Periactin	Tylenol Flu Night Time
Contac	Phenergan	Tylenol PM
	Pyribenzamine	Vicks 44M

IF POSSIBLE, PLEASE DISCONTINUE ANY ANTI-DEPRESSANTS **24 HOURS** prior.

ASTELIN, ASTEPRO AND PATANASE NASAL SPRAYS and BENADRYL LIQUID & TABLETS (DIPHENHYDRAMINE) must be discontinued for **2 DAYS**.

ATARAX, HYDROXYZINE, VISTARIL, ZYRTEC, ZYRTEC-D, CETIRIZINE and XYZAL (Levocetirizine) should be discontinued for **7 DAYS**.

***There is no need to discontinue Corticosteroid Nose Sprays, (Flonase, Nasacort, Nasarel, Nasonex, Rhinocort, Veramyst, Fluticasone, etc.).***